



Erie County  **BRFSS**
Behavioral Risk Factor Surveillance Survey

**Behavioral Health Risks of Erie County Adults
2011**

Table of Contents

2011 Erie County Behavioral Risk Factor Surveillance Survey

Section 1: Health Status

- 1.1 [Percent of adults with fair or poor health](#)

Section 2: Health-Related Quality of Life

- 2.1 [Percent of adults with physical health not good 1+ days in past month](#)
- 2.2 [Percent of adults with mental health not good 1+ days in past month](#)
- 2.3 [Percent of adults who poor physical/mental health prevented usual activity 1+ days in past month](#)

Section 3: Health Care Access

- 3.1 [Percent of adults with no health insurance, age 18-64](#)
- 3.2 [Percent of adults who do not have a personal health care provider](#)
- 3.3 [Percent of adults who needed to see a doctor in past year but could not because of cost](#)
- 3.4 [Percent of adults who last visited doctor for routine checkup within past 2 years](#)

Section 4: Hypertension Awareness

- 4.1 [Percent of adults who ever been told they have high blood pressure](#)
- 4.2 [Percent of adults who are taking medication for high blood pressure](#)

Section 5: Cholesterol Awareness

- 5.1 [Percent of adults who have ever had blood cholesterol checked](#)
- 5.2 [Percent of adults who had blood cholesterol checked in past 5 years](#)
- 5.3 [Percent of adults who have ever been told they have high blood cholesterol](#)

Section 6: Chronic Health Conditions

- 6.1 [Percent of adults who have ever been told they had a heart attack, age 35+](#)
- 6.2 [Percent of adults who have ever been told they had heart disease, age 35+](#)
- 6.3 [Percent of adults who have ever been told they had a stroke, age 35+](#)
- 6.4 [Percent of adults who have ever been told they had asthma](#)
- 6.5 [Percent of adults who currently have asthma](#)
- 6.6 [Percent of adults who have ever been told they had skin cancer](#)
- 6.7 [Percent of adults who have ever been told they had cancer other than skin cancer](#)
- 6.8 [Percent of adults who have ever been told they have COPD, emphysema, or chronic bronchitis](#)
- 6.9 [Percent of adults who have ever been told they have some form of arthritis, rheumatoid arthritis, gout,](#)
- 6.10 [Percent of adults who have ever been told they have a depressive disorder](#)
- 6.11 [Percent of adults who have ever been told they have kidney disease](#)
- 6.12 [Percent of adults who have ever been told they have vision impairment, even when wearing glasses](#)
- 6.13 [Percent of adults who have ever been told they have diabetes](#)

Section 7: Tobacco Use

- 7.1 [Percent of adults who smoked 100+ cigarettes in their lifetime](#)
- 7.2 [Percent of adults who are current smokers](#)
- 7.2 [Percent of adults who are former smokers](#)
- 7.3 [Percent of adults who are smokers who have quit smoking 1+ days in past year](#)
- 7.4 [Percent of adults who last smoked in the past year](#)
- 7.5 [Percent of adults who currently use smokeless tobacco such as chewing tobacco, snuff, or snuss](#)

Section 8: Weight

- 8.11 [Percent of adults who are overweight \(not obese\)](#)
- 8.11 [Percent of adults who are overweight \(including obese\)](#)
- 8.11 [Percent of adults who are obese](#)

Table of Contents

2011 Erie County Behavioral Risk Factor Surveillance Survey

Section 9: Nutrition

9.1-9.5 [Percent of adults who eat fruits and vegetables 5+ times per day](#)

Section 10: Exercise (Physical Activity)

10.1 [Percent of adults with no leisure time physical activity in past month](#)

10.2 [Percent of adults that participated in each activity* \(Out of adults that participated in leisure time activity\)](#)

10.3 [Average number of times spent exercising per week \(of adults who exercised in past month\)](#)

10.4 [Average number of minutes per week spent exercising \(of adults that exercised in the past month\)](#)

10.5 [Average number of times spent strengthening muscles per week \(of adults that did muscle strength activity\)](#)

Section 11: Disability

11.1 [Percent of adults with health problems that require use of special equipment](#)

Section 12: Arthritis Burden

12.1 [Percent of adults who have arthritis or joint pain that limits activity](#)

Section 13: Seatbelt Use

13.1 [Percent of adults who always wear a seat belt when they drive or ride in a car](#)

Section 14: Immunization

14.1 [Percent of adults with seasonal flu shot or flu vaccine that was sprayed up nose in past year, all ages](#)

14.1 [Percent of adults with seasonal flu shot or flu vaccine that was sprayed up nose in past year, age 50+](#)

14.1 [Percent of adults with seasonal flu shot or flu vaccine that was sprayed up nose in past year, age 65+](#)

14.3 [Place where received last flu shot/nasal vaccine spray](#)

14.4 [Percent of adults who ever had a pneumonia vaccination, age 65+](#)

Section 15: Alcohol Consumption

15.1 [Alcohol Consumption: Percent of adults who participated in Chronic Drinking in the past month](#)

15.2 [Alcohol Consumption: Percent of adults who are heavy drinkers](#)

15.3 [Percent of adults who are binge drinkers](#)

Section 16: HIV/AIDS

16.1 [Percent of adults ever tested for HIV, age 18-64](#)

16.2 [Percent of adults who practiced risky behavior within the past year that could result in HIV infection](#)

Module 1: Pre-Diabetes

1.1 [Percent of adults who have ever been told they have pre-diabetes](#)

Module 7: Inadequate Sleep

7.1 [Percent of adults who sleep less than 7 hours in a 24 hour period](#)

Module Erie HC: Health Care Provider

1 [Place where usually go when sick](#)

2 [Where went for care when sick in past 12 months](#)

Module 12: Women's Health

12.1 [Percent of women who had a mammogram in the past year, women age 40+](#)

12.1 [Percent of women who never had a mammogram, women age 40+](#)

12.2 [Percent of women who ever had a clinical breast exam, women age 40+](#)

12.2 [Percent of women who had a clinical breast exam in the past year, women age 40+](#)

12.3 [Percent of women who had a pap test in the past year, women age 18+](#)

Module 13: Prostate Cancer Screening

13.1 [Percent of men who had a prostate-specific antigen test in the past year, men age 40+](#)

13.2 [Percent of men who had a digital rectal exam in the past year, men age 40+](#)

13.3 [Percent of men who have ever been told they had prostate cancer, mean age 40+](#)

Table of Contents

2011 Erie County Behavioral Risk Factor Surveillance Survey

Module 14: Colorectal Cancer Screening

- 14.1 [Percent of adults who had a blood stool test in the past 2 years, age 50+](#)
- 14.2 [Percent of adults who had a sigmoidoscopy or colonoscopy in the past 5 years, age 50+](#)

Module 23: General Preparedness

- 23.1 [Percent of adults who feel household is well prepared to handle a large scale disaster or emergency](#)

Module 28: Social Context

- 28.1 [Percent of adults who were worried or stressed about having enough money](#)
- 28.2 [Percent of adults who needed prescribed medication in past 12 months but did not get it due to cost](#)

Module Erie Tobacco: Tobacco Use

- 1 [Percent of adult smokers who smoke 1+ packs of cigarettes \(20 or more cigarettes\) per day](#)

Module 16: Secondhand Smoke

- 16.1 [Percent of adults where smoking is never allowed inside the house](#)

Module 26: Anxiety and Mental Health

- 26.1 [Percent of adults who have ever been told have an anxiety disorder](#)

Module Erie Alcohol: Alcohol Consumption

- 1 [Percent of adults who drove 1 or more times when they perhaps had too much to drink in the past 30 d](#)

Module Erie Nutri: Nutrition

- 2 [Percent of adults who have 2 or more servings of dairy daily](#)
- 2 [Percent of adults who have 1 or more servings of grains daily](#)
- 3 [Average number of sugar sweetened beverages per day](#)

Module Erie Oral: Oral Health

- 1 [Percent of adults who visited a dentist within the past year](#)

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q1.1: Percent of adults with fair or poor health

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	14%	12%	16%	17%	2.1%	15%	19%	
Male	11%	8%	15%	18%	3.1%	15%	21%	***
Female	16%	14%	19%	16%	2.9%	13%	19%	
18-29	5%	2%	12%	11%	3.7%	8%	15%	
30-44	8%	5%	12%	12%	3.9%	8%	16%	
45-64	17%	14%	21%	19%	3.7%	15%	23%	
65+	25%	19%	31%	26%	5.7%	20%	31%	
< High School	40%	27%	53%	37%	10.0%	27%	47%	
High School	15%	12%	19%	23%	3.9%	19%	27%	***
Some College	10%	7%	15%	19%	4.1%	15%	23%	***
College Graduate	7%	5%	10%	5%	2.5%	2%	7%	
<\$25,000	26%	21%	32%	32%	5.2%	27%	37%	
\$25,000 to \$49,999	14%	10%	19%	11%	3.8%	7%	15%	
\$50,000+	5%	3%	7%	6%	2.3%	4%	8%	
White, non-Hispanic	14%	12%	16%	16%	2.1%	13%	18%	
Black, non-Hispanic	NA			25%	12.0%	12%	37%	
Hispanic	NA			41%	17.9%	23%	59%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q2.1: Percent of adults with physical health not good 1+ days in past month

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	37%	34%	41%	36%	2.7%	34%	39%	
Male	31%	26%	37%	35%	3.9%	31%	39%	
Female	43%	38%	47%	38%	3.9%	34%	42%	
18-29	35%	25%	47%	37%	5.8%	31%	43%	
30-44	34%	28%	41%	38%	5.8%	32%	43%	
45-64	37%	32%	42%	38%	4.6%	33%	42%	
65+	44%	37%	50%	32%	6.2%	26%	38%	
< High School	45%	32%	59%	50%	10.7%	40%	61%	
High School	38%	33%	43%	40%	4.7%	36%	45%	
Some College	38%	31%	45%	41%	5.2%	35%	46%	
College Graduate	33%	28%	39%	24%	4.9%	19%	29%	
<\$25,000	52%	44%	59%	47%	5.6%	41%	53%	
\$25,000 to \$49,999	36%	30%	42%	31%	5.5%	26%	37%	
\$50,000+	30%	25%	35%	32%	4.6%	27%	37%	
White, non-Hispanic	37%	33%	40%	36%	2.9%	34%	39%	
Black, non-Hispanic	NA			32%	13.3%	19%	45%	
Hispanic	NA			43%	18.3%	25%	61%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q2.2: Percent of adults with mental health not good 1+ days in past month

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	35%	31%	38%	33%	2.7%	30%	35%	
Male	29%	24%	35%	25%	3.5%	21%	28%	
Female	40%	36%	44%	40%	3.9%	36%	44%	
18-29	52%	41%	63%	42%	5.9%	36%	48%	
30-44	38%	31%	44%	37%	5.8%	31%	43%	
45-64	32%	28%	37%	32%	4.5%	28%	37%	
65+	16%	12%	22%	17%	4.9%	12%	22%	
< High School	49%	35%	62%	47%	10.7%	36%	57%	
High School	30%	25%	35%	35%	4.5%	30%	39%	
Some College	41%	33%	49%	37%	5.1%	32%	42%	
College Graduate	32%	26%	38%	22%	4.8%	17%	27%	
<\$25,000	40%	33%	47%	45%	5.6%	39%	51%	
\$25,000 to \$49,999	38%	31%	45%	27%	5.3%	22%	32%	
\$50,000+	29%	24%	35%	32%	4.6%	27%	37%	
White, non-Hispanic	34%	30%	37%	32%	2.8%	29%	35%	
Black, non-Hispanic	NA			44%	14.3%	29%	58%	
Hispanic	NA			41%	17.9%	23%	59%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q2.3: Percent of adults in which poor physical/mental health prevented usual activity 1+ days in past month

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			21%	2.3%	19%	23%	
Male	NA			19%	3.2%	16%	22%	
Female	NA			23%	3.3%	20%	27%	
18-29	NA			22%	5.0%	17%	27%	
30-44	NA			24%	5.1%	19%	29%	
45-64	NA			24%	4.0%	19%	28%	
65+	NA			11%	4.2%	7%	15%	
< High School	NA			21%	8.4%	13%	30%	
High School	NA			23%	3.9%	19%	27%	
Some College	NA			26%	4.6%	22%	31%	
College Graduate	NA			13%	4.0%	9%	17%	
<\$25,000	NA			35%	5.3%	30%	40%	
\$25,000 to \$49,999	NA			17%	4.5%	13%	22%	
\$50,000+	NA			16%	3.6%	13%	20%	
White, non-Hispanic	NA			21%	2.4%	19%	23%	
Black, non-Hispanic	NA			28%	12.8%	15%	40%	
Hispanic	NA			21%	15.2%	6%	37%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q3.1: Percent of adults with no health insurance, age 18-64

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	17%	13%	21%	13%	2.1%	11%	15%	
Male	22%	17%	29%	13%	3.0%	10%	16%	***
Female	11%	7%	15%	13%	2.9%	10%	15%	
18-29	33%	22%	45%	19%	4.6%	14%	24%	
30-44	15%	10%	21%	15%	4.3%	11%	19%	
45-64	9%	6%	12%	8%	2.5%	5%	10%	
65+	NA			NA				
< High School	NSR			24%	10.3%	14%	35%	
High School	19%	14%	26%	18%	4.1%	14%	22%	
Some College	15%	9%	23%	14%	3.9%	10%	17%	
College Graduate	6%	3%	12%	6%	2.9%	3%	8%	
<\$25,000	35%	25%	45%	25%	5.5%	19%	30%	
\$25,000 to \$49,999	16%	11%	24%	14%	4.5%	9%	18%	
\$50,000+	5%	3%	11%	3%	1.8%	1%	5%	
White, non-Hispanic	16%	13%	20%	12%	2.1%	9%	14%	
Black, non-Hispanic	NA			29%	12.7%	16%	41%	
Hispanic	NA			29%	16.7%	12%	45%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q3.2: Percent of adults who do not have a personal health care provider

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	11%	9%	14%	10%	1.7%	8%	12%	
Male	15%	11%	20%	14%	2.8%	11%	17%	
Female	8%	6%	11%	6%	1.9%	4%	8%	
18-29	25%	17%	36%	19%	4.7%	14%	24%	
30-44	12%	8%	17%	13%	4.1%	9%	17%	
45-64	7%	5%	10%	5%	2.1%	3%	8%	
65+	5%	3%	8%	3%	2.3%	1%	5%	
< High School	NSR			17%	7.6%	9%	24%	
High School	11%	8%	16%	10%	2.8%	7%	13%	
Some College	12%	8%	18%	13%	3.5%	9%	17%	
College Graduate	8%	5%	12%	6%	2.7%	3%	8%	
<\$25,000	20%	14%	27%	14%	3.9%	10%	18%	
\$25,000 to \$49,999	10%	6%	15%	9%	3.5%	6%	13%	
\$50,000+	7%	5%	11%	6%	2.3%	4%	8%	
White, non-Hispanic	11%	9%	14%	9%	1.7%	8%	11%	
Black, non-Hispanic	NA			17%	10.7%	6%	28%	
Hispanic	NA			21%	14.7%	6%	35%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q3.3: Percent of adults who needed to see a doctor in past year but could not because of cost

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	9%	7%	11%	13%	1.9%	11%	15%	***
Male	8%	5%	11%	10%	2.5%	8%	13%	
Female	10%	8%	13%	15%	2.8%	12%	18%	
18-29	13%	7%	21%	19%	4.7%	15%	24%	
30-44	13%	9%	18%	19%	4.6%	14%	23%	
45-64	7%	5%	10%	11%	2.9%	8%	14%	
65+	4%	2%	7%	2%	2.0%	0%	4%	
< High School	NSR			34%	9.7%	24%	43%	
High School	9%	7%	13%	15%	3.3%	12%	19%	
Some College	8%	5%	12%	15%	3.7%	11%	19%	
College Graduate	6%	4%	10%	5%	2.5%	2%	7%	
<\$25,000	15%	11%	21%	23%	4.7%	18%	27%	
\$25,000 to \$49,999	10%	7%	15%	12%	4.0%	8%	16%	
\$50,000+	4%	2%	7%	6%	2.4%	4%	9%	
White, non-Hispanic	8%	7%	11%	12%	1.9%	10%	14%	
Black, non-Hispanic	NA			20%	11.3%	9%	32%	
Hispanic	NA			31%	16.8%	14%	48%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q3.4: Percent of adults who last visited doctor for routine checkup within past 2 years

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	81%	77%	83%	86%	2.0%	84%	88%	***
Male	77%	72%	82%	85%	2.9%	82%	88%	***
Female	84%	80%	87%	88%	2.6%	85%	90%	
18-29	70%	58%	79%	78%	4.9%	73%	83%	
30-44	71%	65%	77%	83%	4.5%	78%	87%	***
45-64	86%	82%	89%	89%	3.0%	86%	92%	
65+	94%	90%	96%	95%	2.8%	93%	98%	
< High School	NSR			99%	8.3%	91%	100%	
High School	82%	77%	87%	94%	3.4%	91%	98%	***
Some College	85%	78%	90%	86%	3.6%	83%	90%	
College Graduate	77%	71%	82%	70%	3.5%	67%	74%	
<\$25,000	79%	72%	85%	89%	3.6%	85%	92%	***
\$25,000 to \$49,999	85%	79%	89%	72%	4.6%	67%	76%	***
\$50,000+	78%	73%	83%	98%	3.2%	95%	100%	***
White, non-Hispanic	80%	77%	83%	86%	2.0%	84%	88%	***
Black, non-Hispanic	NA			94%	6.7%	87%	100%	
Hispanic	NA			75%	16.0%	59%	91%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q4.1: Percent of adults who ever been told they have high blood pressure

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	28%	25%	30%	31%	2.6%	29%	34%	
Male	28%	23%	32%	36%	3.9%	32%	40%	***
Female	27%	24%	31%	27%	3.5%	24%	31%	
18-29	9%	4%	17%	9%	3.3%	5%	12%	
30-44	14%	10%	20%	22%	4.9%	17%	27%	
45-64	30%	26%	35%	40%	4.7%	35%	44%	***
65+	60%	53%	66%	55%	6.5%	49%	62%	
< High School	38%	26%	52%	41%	10.3%	31%	51%	
High School	32%	27%	37%	37%	4.6%	33%	42%	
Some College	21%	16%	27%	31%	4.8%	26%	36%	
College Graduate	24%	19%	29%	22%	4.8%	18%	27%	
<\$25,000	35%	29%	41%	41%	5.5%	35%	46%	
\$25,000 to \$49,999	27%	22%	33%	25%	5.1%	20%	30%	
\$50,000+	21%	17%	25%	29%	4.5%	25%	34%	***
White, non-Hispanic	28%	25%	31%	31%	2.7%	29%	34%	
Black, non-Hispanic	NA			39%	13.6%	25%	52%	
Hispanic	NA			24%	15.6%	9%	40%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q4.2: Percent of adults who are taking medication for high blood pressure

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	80%	74%	84%	79%	4.2%	74%	83%	
Male	73%	64%	80%	77%	5.8%	71%	83%	
Female	86%	78%	91%	81%	5.9%	75%	87%	
18-29	NA			26%	17.9%	8%	44%	
30-44	NA			55%	12.8%	42%	68%	
45-64	88%	82%	93%	85%	5.5%	79%	90%	
65+	92%	86%	95%	92%	4.8%	87%	97%	
< High School	NSR			81%	17.3%	64%	99%	
High School	83%	75%	89%	90%	6.4%	83%	96%	
Some College	NSR			75%	8.2%	67%	83%	
College Graduate	82%	71%	89%	65%	7.9%	57%	73%	
<\$25,000	83%	72%	90%	76%	7.5%	69%	83%	
\$25,000 to \$49,999	74%	63%	83%	65%	9.8%	55%	75%	
\$50,000+	80%	69%	88%	92%	7.2%	85%	100%	
White, non-Hispanic	80%	73%	84%	80%	4.2%	76%	84%	
Black, non-Hispanic	NA			63%	21.7%	42%	85%	
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q5.1: Percent of adults who have ever had blood cholesterol checked

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	80%	76%	83%	79%	2.3%	77%	82%	
Male	76%	70%	81%	80%	3.4%	76%	83%	
Female	84%	80%	87%	79%	3.3%	76%	82%	
18-29	40%	29%	51%	42%	6.1%	36%	48%	
30-44	78%	72%	83%	76%	5.2%	71%	81%	
45-64	94%	92%	96%	95%	2.2%	92%	97%	
65+	97%	94%	98%	96%	2.6%	93%	99%	
< High School	NSR			85%	10.3%	75%	95%	
High School	81%	76%	86%	87%	4.0%	83%	91%	
Some College	74%	65%	82%	75%	4.6%	70%	79%	
College Graduate	86%	80%	90%	69%	3.7%	66%	73%	***
<\$25,000	68%	59%	75%	75%	4.9%	70%	80%	
\$25,000 to \$49,999	82%	75%	88%	69%	4.9%	64%	74%	***
\$50,000+	87%	81%	91%	96%	3.4%	93%	100%	***
White, non-Hispanic	81%	78%	85%	80%	2.4%	78%	83%	
Black, non-Hispanic	NA			75%	12.3%	63%	87%	
Hispanic	NA			52%	18.8%	33%	71%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q5.2: Percent of adults who had blood cholesterol checked in past 5 years

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	76%	72%	79%	76%	2.5%	73%	78%	
Male	72%	66%	77%	75%	3.6%	71%	79%	
Female	80%	75%	83%	76%	3.5%	72%	79%	
18-29	36%	26%	47%	38%	6.1%	32%	44%	
30-44	72%	65%	77%	70%	5.6%	64%	76%	
45-64	90%	87%	93%	91%	2.7%	88%	94%	
65+	94%	90%	97%	96%	2.8%	93%	98%	
< High School	NSR			82%	10.8%	71%	92%	
High School	78%	72%	83%	83%	4.2%	79%	87%	
Some College	71%	62%	79%	71%	4.8%	66%	76%	
College Graduate	78%	72%	83%	65%	4.3%	61%	70%	***
<\$25,000	63%	55%	71%	72%	5.2%	66%	77%	
\$25,000 to \$49,999	79%	72%	85%	65%	5.2%	60%	70%	***
\$50,000+	81%	76%	86%	91%	3.9%	88%	95%	***
White, non-Hispanic	77%	73%	80%	76%	2.6%	74%	79%	
Black, non-Hispanic	NA			73%	12.6%	60%	85%	
Hispanic	NA			52%	18.8%	33%	71%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q5.3: Percent of adults who have ever been told they have high blood cholesterol

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	38%	35%	42%	39%	3.2%	36%	42%	
Male	39%	34%	45%	41%	4.6%	36%	45%	
Female	38%	33%	42%	38%	4.4%	33%	42%	
18-29	NA			15%	6.8%	8%	22%	
30-44	20%	15%	26%	25%	6.1%	19%	31%	
45-64	45%	40%	50%	46%	4.9%	41%	51%	
65+	58%	51%	64%	51%	6.9%	44%	58%	
< High School	NSR			48%	13.3%	35%	62%	
High School	40%	35%	46%	48%	5.5%	42%	53%	
Some College	36%	30%	44%	41%	6.0%	35%	47%	
College Graduate	37%	31%	44%	26%	5.5%	20%	31%	***
<\$25,000	49%	42%	57%	45%	6.6%	39%	52%	
\$25,000 to \$49,999	38%	32%	44%	36%	6.5%	29%	43%	
\$50,000+	34%	28%	39%	35%	5.2%	30%	40%	
White, non-Hispanic	38%	35%	42%	39%	3.3%	36%	43%	
Black, non-Hispanic	NA			33%	15.4%	18%	49%	
Hispanic	NA			36%	25.1%	11%	61%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.1: Percent of adults who have ever been told they had a heart attack, age 35+

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	5%	4%	7%	6%	1.6%	4%	7%	
Male	6%	4%	9%	7%	2.7%	5%	10%	
Female	4%	3%	6%	4%	1.9%	2%	6%	
18-34	NA			NA				
35-44	NA			2%	2.2%	0%	4%	
45-64	NA			5%	2.0%	3%	6%	
65+	12%	8%	17%	10%	3.9%	6%	13%	
< High School	NSR			29%	11.8%	17%	41%	
High School	4%	3%	7%	5%	2.4%	3%	8%	
Some College	5%	2%	9%	6%	3.1%	3%	9%	
College Graduate	5%	3%	8%	2%	2.1%	0%	4%	
<\$25,000	8%	5%	13%	11%	4.3%	7%	15%	
\$25,000 to \$49,999	5%	3%	8%	3%	2.4%	0%	5%	
\$50,000+	3%	2%	5%	2%	1.5%	0%	3%	
White, non-Hispanic	5%	4%	7%	5%	1.6%	4%	7%	
Black, non-Hispanic	NA			0%	NCI			
Hispanic	NA			9%	17.0%	0%	26%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.2: Percent of adults who have ever been told they had heart disease, age 35+

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	8%	6%	10%	7%	1.8%	5%	9%	
Male	9%	6%	12%	10%	3.0%	7%	13%	
Female	7%	5%	10%	5%	2.1%	3%	7%	
18-34	NA			NA				
35-44	NA			2%	2.2%	0%	4%	
45-64	NA			6%	2.3%	4%	9%	
65+	18%	13%	24%	12%	4.2%	7%	16%	
< High School	NSR			19%	10.2%	9%	29%	
High School	7%	5%	10%	7%	2.8%	4%	10%	
Some College	7%	4%	11%	6%	3.2%	3%	9%	
College Graduate	5%	3%	9%	6%	3.6%	2%	10%	
<\$25,000	11%	7%	16%	9%	4.0%	5%	13%	
\$25,000 to \$49,999	9%	6%	13%	3%	2.7%	0%	6%	***
\$50,000+	5%	3%	8%	8%	3.2%	5%	11%	
White, non-Hispanic	8%	6%	10%	7%	1.8%	5%	9%	
Black, non-Hispanic	NA			4%	6.9%	0%	11%	
Hispanic	NA			9%	17.0%	0%	26%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.3: Percent of adults who have ever been told they had a stroke, age 35+

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	4%	3%	6%	5%	1.6%	4%	7%	
Male	3%	2%	6%	5%	2.2%	3%	7%	
Female	5%	3%	7%	6%	2.2%	3%	8%	
18-34	NA			NA				
35-44	NA			2%	2.6%	0%	5%	
45-64	NA			4%	1.9%	2%	6%	
65+	10%	6%	15%	9%	3.8%	5%	13%	
< High School	NSR			20%	10.6%	9%	31%	
High School	5%	3%	7%	6%	2.5%	3%	8%	
Some College	3%	1%	6%	6%	3.2%	3%	9%	
College Graduate	2%	1%	4%	1%	1.7%	0%	3%	
<\$25,000	8%	5%	13%	7%	3.5%	3%	11%	
\$25,000 to \$49,999	6%	4%	10%	5%	3.2%	1%	8%	
\$50,000+	NA			1%	1.3%	0%	3%	
White, non-Hispanic	4%	3%	6%	4%	1.5%	3%	6%	
Black, non-Hispanic	NA			21%	15.2%	6%	37%	
Hispanic	NA			0%	NCI			

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.4: Percent of adults who have ever been told they had asthma

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	8%	7%	10%	12%	1.8%	10%	14%	***
Male	5%	3%	8%	10%	2.4%	7%	12%	
Female	11%	9%	14%	14%	2.8%	12%	17%	
18-29	11%	6%	20%	17%	4.5%	13%	22%	
30-44	8%	5%	11%	14%	4.1%	10%	18%	
45-64	9%	7%	12%	10%	2.8%	7%	13%	
65+	6%	3%	11%	8%	3.6%	5%	12%	
< High School	11%	5%	23%	19%	7.8%	11%	27%	
High School	6%	4%	10%	13%	3.1%	10%	16%	***
Some College	11%	8%	17%	14%	3.6%	10%	17%	
College Graduate	8%	5%	12%	8%	3.2%	5%	11%	
<\$25,000	11%	7%	17%	16%	4.1%	12%	20%	
\$25,000 to \$49,999	7%	4%	10%	11%	3.7%	7%	14%	
\$50,000+	8%	5%	11%	11%	3.0%	8%	14%	
White, non-Hispanic	8%	6%	10%	12%	1.9%	10%	14%	***
Black, non-Hispanic	NA			6%	6.7%	0%	13%	
Hispanic	NA			24%	15.6%	9%	40%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.5: Percent of adults who currently have asthma

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	6%	5%	8%	8%	1.5%	6%	9%	
Male	4%	2%	7%	6%	2.0%	4%	8%	
Female	8%	6%	11%	10%	2.3%	7%	12%	
18-29	7%	3%	16%	10%	3.6%	7%	14%	
30-44	6%	3%	9%	10%	3.6%	6%	13%	
45-64	6%	4%	9%	7%	2.4%	4%	9%	
65+	5%	2%	10%	5%	2.7%	2%	7%	
< High School	11%	5%	23%	12%	6.5%	5%	18%	
High School	5%	3%	8%	9%	2.7%	7%	12%	
Some College	8%	5%	12%	7%	2.7%	5%	10%	
College Graduate	5%	3%	8%	5%	2.7%	3%	8%	
<\$25,000	8%	5%	14%	12%	3.7%	9%	16%	
\$25,000 to \$49,999	4%	3%	7%	5%	2.6%	2%	8%	
\$50,000+	6%	4%	9%	8%	2.6%	5%	11%	
White, non-Hispanic	5%	4%	7%	8%	1.6%	6%	9%	
Black, non-Hispanic	NA			6%	6.7%	0%	13%	
Hispanic	NA			18%	14.2%	4%	32%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.6: Percent of adults who have ever been told they had skin cancer

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			5%	1.2%	4%	6%	
Male	NA			4%	1.6%	2%	5%	
Female	NA			6%	1.9%	4%	8%	
18-29	NA			0%	NCI			
30-44	NA			1%	1.0%	0%	2%	
45-64	NA			7%	2.4%	4%	9%	
65+	NA			12%	4.3%	8%	17%	
< High School	NA			12%	6.5%	5%	18%	
High School	NA			5%	2.0%	3%	7%	
Some College	NA			4%	2.0%	2%	6%	
College Graduate	NA			4%	2.2%	1%	6%	
<\$25,000	NA			3%	1.9%	1%	5%	
\$25,000 to \$49,999	NA			4%	2.5%	2%	7%	
\$50,000+	NA			6%	2.4%	4%	9%	
White, non-Hispanic				5%	1.3%	4%	6%	
Black, non-Hispanic	NA			2%	4.1%	0%	6%	
Hispanic	NA			3%	6.6%	0%	10%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.7: Percent of adults who have ever been told they had cancer other than skin cancer

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	NA			8%	1.5%	6%	9%	
Male	NA			5%	1.8%	3%	7%	
Female	NA			10%	2.4%	8%	13%	
18-29	NA			2%	1.4%	0%	3%	
30-44	NA			2%	1.5%	0%	3%	
45-64	NA			9%	2.7%	6%	11%	
65+	NA			22%	5.4%	16%	27%	
< High School	NA			12%	6.5%	5%	18%	
High School	NA			10%	2.8%	8%	13%	
Some College	NA			7%	2.6%	4%	9%	
College Graduate	NA			5%	2.5%	2%	7%	
<\$25,000	NA			12%	3.7%	8%	16%	
\$25,000 to \$49,999	NA			7%	3.1%	4%	10%	
\$50,000+	NA			6%	2.3%	4%	8%	
White, non-Hispanic	NA			8%	1.6%	6%	9%	
Black, non-Hispanic	NA			6%	7.0%	0%	13%	
Hispanic	NA			0%	NCI			

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.8: Percent of adults who have ever been told they have COPD, emphysema, or chronic bronchitis

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	NA			7%	1.5%	6%	9%	
Male	NA			7%	2.1%	5%	9%	
Female	NA			7%	2.0%	5%	9%	
18-29	NA			4%	2.3%	2%	6%	
30-44	NA			2%	1.8%	0%	4%	
45-64	NA			8%	2.6%	5%	10%	
65+	NA			15%	4.7%	10%	20%	
< High School	NA			17%	7.6%	9%	24%	
High School	NA			13%	3.1%	10%	16%	
Some College	NA			5%	2.3%	3%	7%	
College Graduate	NA			1%	1.4%	0%	3%	
<\$25,000	NA			13%	3.8%	10%	17%	
\$25,000 to \$49,999	NA			4%	2.5%	2%	7%	
\$50,000+	NA			5%	2.1%	3%	7%	
White, non-Hispanic	NA			7%	1.5%	5%	8%	
Black, non-Hispanic	NA			2%	4.1%	0%	6%	
Hispanic	NA			21%	14.7%	6%	35%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.9: Percent of adults who have ever been told they have some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	30%	28%	34%	30%	2.6%	28%	33%	
Male	26%	22%	30%	25%	3.6%	22%	29%	
Female	35%	31%	39%	35%	3.8%	31%	39%	
18-29	5%	2%	12%	4%	2.3%	2%	6%	
30-44	15%	10%	20%	15%	4.3%	11%	19%	
45-64	37%	32%	42%	41%	4.7%	37%	46%	
65+	66%	59%	71%	61%	6.4%	54%	67%	
< High School	35%	23%	48%	48%	10.6%	38%	59%	
High School	34%	30%	39%	41%	4.7%	37%	46%	
Some College	28%	22%	34%	30%	4.8%	25%	34%	
College Graduate	26%	21%	31%	16%	4.3%	12%	20%	***
<\$25,000	39%	33%	46%	40%	5.5%	34%	45%	
\$25,000 to \$49,999	33%	28%	39%	24%	5.1%	19%	30%	
\$50,000+	22%	18%	26%	23%	4.1%	19%	27%	
White, non-Hispanic	31%	28%	34%	31%	2.7%	28%	34%	
Black, non-Hispanic	NA			20%	11.3%	9%	32%	
Hispanic	NA			17%	13.3%	3%	30%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.10: Percent of adults who have ever been told they have a depressive disorder

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			19%	2.3%	17%	22%	
Male	NA			17%	3.0%	13%	20%	
Female	NA			22%	3.3%	19%	25%	
18-29	NA			24%	5.0%	19%	29%	
30-44	NA			16%	4.4%	12%	20%	
45-64	NA			22%	4.0%	18%	26%	
65+	NA			13%	4.4%	8%	17%	
< High School	NA			28%	9.1%	19%	37%	
High School	NA			22%	3.8%	18%	26%	
Some College	NA			22%	4.4%	18%	27%	
College Graduate	NA			12%	3.8%	8%	16%	
<\$25,000	NA			31%	5.2%	25%	36%	
\$25,000 to \$49,999	NA			15%	4.3%	11%	20%	
\$50,000+	NA			14%	3.4%	10%	17%	
White, non-Hispanic	NA			20%	2.3%	17%	22%	
Black, non-Hispanic	NA			14%	9.8%	4%	24%	
Hispanic	NA			29%	16.7%	12%	45%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.11: Percent of adults who have ever been told they have kidney disease

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			2%	0.8%	1%	3%	
Male	NA			3%	1.3%	1%	4%	
Female	NA			2%	1.0%	1%	3%	
18-29	NA			0%	NCI			
30-44	NA			2%	1.5%	0%	3%	
45-64	NA			3%	1.6%	1%	4%	
65+	NA			5%	2.8%	2%	7%	
< High School	NA			7%	5.2%	2%	13%	
High School	NA			2%	1.4%	1%	4%	
Some College	NA			3%	1.6%	1%	4%	
College Graduate	NA			0%	0.8%	0%	1%	
<\$25,000	NA			3%	2.0%	1%	5%	
\$25,000 to \$49,999	NA			2%	1.6%	0%	4%	
\$50,000+	NA			1%	0.7%	0%	1%	
White, non-Hispanic				2%	0.8%	1%	3%	
Black, non-Hispanic	NA			2%	4.1%	0%	6%	
Hispanic	NA			7%	9.2%	0%	16%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.12: Percent of adults who have ever been told they have vision impairment, even when wearing glasses

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			14%	2.0%	12%	16%	
Male	NA			11%	2.6%	9%	14%	
Female	NA			17%	3.0%	14%	20%	
18-29	NA			10%	3.5%	6%	13%	
30-44	NA			10%	3.6%	7%	14%	
45-64	NA			13%	3.2%	9%	16%	
65+	NA			28%	5.9%	22%	34%	
< High School	NA			21%	8.3%	13%	29%	
High School	NA			16%	3.4%	12%	19%	
Some College	NA			16%	3.8%	12%	19%	
College Graduate	NA			10%	3.5%	6%	14%	
<\$25,000	NA			23%	4.7%	18%	28%	
\$25,000 to \$49,999	NA			10%	3.6%	7%	14%	
\$50,000+	NA			11%	3.1%	8%	14%	
White, non-Hispanic				14%	2.1%	12%	16%	
Black, non-Hispanic	NA			10%	8.6%	2%	19%	
Hispanic	NA			10%	11.1%	0%	21%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q6.13: Percent of adults who have ever been told they have diabetes

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	8%	7%	10%	10%	1.7%	8%	12%	
Male	8%	6%	11%	10%	2.5%	8%	13%	
Female	9%	7%	11%	10%	2.4%	8%	12%	
18-29	0%	NCI		1%	1.0%	0%	2%	
30-44	2%	1%	4%	6%	2.7%	3%	8%	
45-64	11%	8%	14%	14%	3.3%	10%	17%	
65+	20%	15%	26%	21%	5.4%	15%	26%	
< High School	19%	11%	32%	12%	6.4%	5%	18%	
High School	9%	7%	12%	14%	3.2%	10%	17%	
Some College	7%	4%	11%	11%	3.3%	8%	14%	
College Graduate	6%	4%	9%	5%	2.6%	3%	8%	
<\$25,000	13%	9%	17%	15%	4.0%	11%	19%	
\$25,000 to \$49,999	7%	4%	10%	7%	3.0%	4%	10%	
\$50,000+	6%	4%	8%	8%	2.6%	5%	11%	
White, non-Hispanic	8%	6%	10%	10%	1.8%	9%	12%	
Black, non-Hispanic	NA			8%	7.7%	1%	16%	
Hispanic	NA			7%	9.2%	0%	16%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q7.1: Percent of adults who smoked 100+ cigarettes in their lifetime

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	52%	48%	55%	50%	2.8%	47%	53%	
Male	57%	52%	63%	52%	4.1%	48%	56%	
Female	47%	43%	51%	48%	4.0%	44%	52%	
18-29	46%	35%	57%	39%	5.8%	33%	45%	
30-44	52%	45%	59%	46%	6.0%	40%	52%	
45-64	56%	51%	60%	56%	4.7%	51%	60%	
65+	54%	47%	60%	59%	6.5%	52%	65%	
< High School	NSR			80%	10.2%	69%	90%	
High School	54%	49%	59%	64%	4.8%	60%	69%	
Some College	54%	46%	62%	49%	5.2%	44%	55%	
College Graduate	41%	35%	48%	30%	5.2%	25%	35%	***
<\$25,000	59%	51%	66%	61%	5.5%	55%	66%	
\$25,000 to \$49,999	57%	50%	63%	48%	5.7%	43%	54%	
\$50,000+	44%	38%	49%	44%	5.0%	39%	49%	
White, non-Hispanic	52%	48%	55%	50%	3.0%	47%	53%	
Black, non-Hispanic	NA			55%	13.9%	41%	69%	
Hispanic	NA			52%	18.2%	34%	70%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q7.2: Percent of adults who are current smokers

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	26%	23%	29%	23%	2.4%	21%	25%	
Male	28%	23%	34%	22%	3.4%	19%	26%	
Female	24%	21%	28%	24%	3.4%	20%	27%	
18-29	35%	25%	47%	27%	5.3%	22%	32%	
30-44	34%	28%	40%	28%	5.4%	23%	33%	
45-64	26%	22%	30%	23%	4.0%	19%	27%	
65+	8%	5%	13%	10%	4.0%	6%	14%	
< High School	47%	33%	60%	42%	10.3%	32%	52%	
High School	32%	27%	38%	30%	4.3%	26%	34%	
Some College	27%	21%	34%	24%	4.5%	20%	29%	
College Graduate	10%	7%	15%	10%	3.6%	7%	14%	
<\$25,000	37%	30%	45%	37%	5.4%	31%	42%	
\$25,000 to \$49,999	32%	26%	38%	21%	4.9%	16%	26%	***
\$50,000+	16%	13%	21%	15%	3.5%	12%	19%	
White, non-Hispanic	25%	22%	29%	22%	2.4%	19%	24%	
Black, non-Hispanic	NA			35%	13.3%	21%	48%	
Hispanic	NA			31%	16.8%	14%	48%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q7.2: Percent of adults who are former smokers

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	26%	23%	29%	27%	2.5%	25%	30%	
Male	29%	25%	34%	30%	3.7%	26%	34%	
Female	23%	20%	27%	25%	3.4%	21%	28%	
18-29	10%	6%	19%	12%	3.9%	8%	16%	
30-44	18%	14%	24%	18%	4.6%	13%	22%	
45-64	30%	25%	34%	32%	4.5%	28%	37%	
65+	45%	39%	52%	48%	6.6%	42%	55%	
< High School	24%	15%	36%	38%	10.0%	28%	48%	
High School	22%	18%	26%	34%	4.5%	30%	39%	***
Some College	27%	22%	34%	25%	4.5%	20%	29%	
College Graduate	31%	26%	37%	20%	4.6%	15%	24%	***
<\$25,000	22%	17%	27%	24%	4.8%	19%	29%	
\$25,000 to \$49,999	25%	21%	31%	27%	5.3%	22%	32%	
\$50,000+	27%	23%	32%	29%	4.5%	24%	33%	
White, non-Hispanic	26%	24%	29%	28%	2.6%	25%	31%	
Black, non-Hispanic	NA			20%	11.3%	9%	32%	
Hispanic	NA			21%	14.7%	6%	35%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q7.3: Percent of adults who are smokers who have quit smoking 1+ days in past year

	2007		2011				Sig
	%	Lower Upper	%	C I	Lower	Upper	
Total	56%	NA	57%	5.9%	51%	63%	
Male	54%	NA	62%	8.4%	53%	70%	
Female	59%	NA	52%	8.2%	44%	61%	
18-29	71%	NA	69%	10.4%	59%	80%	
30-44	52%	NA	53%	11.3%	42%	65%	
45-64	50%	NA	52%	9.8%	42%	61%	
65+	54%	NA	48%	20.4%	27%	68%	
< High School	67%	NA	60%	17.9%	42%	78%	
High School	48%	NA	62%	9.1%	53%	71%	
Some College	68%	NA	57%	10.6%	47%	68%	
College Graduate	55%	NA	49%	14.4%	35%	64%	
<\$25,000	57%	NA	57%	9.2%	48%	66%	
\$25,000 to \$49,999	59%	NA	48%	11.8%	36%	60%	
\$50,000+	54%	NA	57%	13.8%	43%	71%	
White, non-Hispanic	56%	NA	56%	6.3%	50%	62%	
Black, non-Hispanic	56%	NA	65%	22.7%	42%	87%	
Hispanic	86%	NA	80%	24.8%	55%	100%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q7.4: Percent of adults who last smoked in the past year

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	11%	7%	17%	13%	3.7%	9%	17%	
Male	13%	7%	22%	10%	4.5%	6%	15%	
Female	9%	5%	17%	16%	5.8%	10%	22%	
18-29	NSR			61%	16.7%	44%	77%	
30-44	NSR			25%	12.0%	12%	37%	
45-64	8%	4%	15%	7%	4.2%	2%	11%	
65+	3%	1%	8%	2%	2.6%	0%	4%	
< High School	NSR			14%	12.3%	2%	26%	
High School	10%	4%	19%	10%	5.1%	5%	15%	
Some College	17%	9%	29%	13%	7.0%	6%	20%	
College Graduate	NSR			15%	8.5%	7%	24%	
<\$25,000	NSR			21%	9.3%	11%	30%	
\$25,000 to \$49,999	10%	5%	22%	12%	7.2%	5%	20%	
\$50,000+	NSR			13%	6.4%	7%	20%	
White, non-Hispanic	10%	6%	15%	12%	3.6%	8%	16%	
Black, non-Hispanic	NA			40%	30.4%	10%	70%	
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q7.5: Percent of adults who currently use smokeless tobacco such as chewing tobacco, snuff, or snus

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	3%	2%	6%	4%	1.1%	3%	5%	
Male	7%	4%	11%	8%	2.2%	6%	11%	
Female	0%	0%	2%	0%	NCI			
18-29	NSR			7%	2.9%	4%	10%	
30-44	6%	3%	11%	4%	2.4%	2%	7%	
45-64	2%	1%	5%	3%	1.6%	1%	5%	
65+	1%	0%	4%	3%	2.3%	1%	5%	
< High School	NSR			0%	NCI			
High School	5%	3%	10%	6%	2.2%	4%	8%	
Some College	2%	0%	5%	6%	2.5%	4%	9%	
College Graduate	1%	1%	4%	1%	1.4%	0%	3%	
<\$25,000	3%	1%	11%	4%	2.2%	2%	6%	
\$25,000 to \$49,999	2%	1%	5%	3%	2.1%	1%	5%	
\$50,000+	5%	2%	9%	5%	2.2%	3%	7%	
White, non-Hispanic	3%	2%	6%	4%	1.2%	3%	5%	
Black, non-Hispanic	NA			0%	NCI			
Hispanic	NA			7%	9.2%	0%	16%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q8.11: Percent of adults who are overweight (not obese)

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			36%	2.8%	33%	39%	
Male	NA			40%	4.0%	36%	44%	
Female	NA			32%	3.8%	28%	36%	
18-29	NA			31%	5.5%	25%	37%	
30-44	NA			29%	5.5%	24%	35%	
45-64	NA			40%	4.8%	35%	45%	
65+	NA			44%	6.7%	37%	51%	
< High School	NA			42%	10.4%	32%	53%	
High School	NA			41%	4.8%	36%	45%	
Some College	NA			34%	5.0%	29%	39%	
College Graduate	NA			30%	5.3%	25%	35%	
<\$25,000	NA			32%	5.2%	26%	37%	
\$25,000 to \$49,999	NA			31%	5.5%	25%	36%	
\$50,000+	NA			45%	5.1%	40%	50%	
White, non-Hispanic	NA			37%	2.9%	34%	40%	
Black, non-Hispanic	NA			23%	11.9%	11%	35%	
Hispanic	NA			31%	16.8%	14%	48%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q8.11: Percent of adults who are overweight (including obese)

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	65%	62%	69%	65%	2.8%	62%	68%	
Male	71%	65%	76%	71%	3.7%	67%	74%	
Female	60%	56%	64%	59%	4.0%	55%	63%	
18-29	53%	42%	65%	53%	6.0%	47%	59%	
30-44	68%	62%	74%	62%	5.9%	56%	67%	
45-64	69%	65%	74%	71%	4.4%	67%	75%	
65+	67%	60%	72%	72%	6.0%	66%	78%	
< High School	65%	51%	77%	67%	10.7%	57%	78%	
High School	70%	64%	75%	73%	4.7%	68%	77%	
Some College	59%	51%	67%	69%	4.9%	64%	74%	
College Graduate	64%	58%	70%	49%	5.3%	44%	54%	***
<\$25,000	67%	59%	73%	64%	5.4%	59%	70%	
\$25,000 to \$49,999	63%	57%	69%	56%	5.6%	50%	62%	
\$50,000+	69%	63%	74%	74%	4.8%	69%	79%	
White, non-Hispanic	65%	61%	68%	65%	2.9%	62%	68%	
Black, non-Hispanic	NA			59%	13.8%	45%	73%	
Hispanic	NA			66%	17.3%	48%	83%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q8.11: Percent of adults who are obese

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	28%	25%	31%	29%	2.6%	26%	31%	
Male	28%	23%	33%	30%	3.8%	26%	34%	
Female	29%	25%	33%	27%	3.6%	23%	31%	
18-29	23%	15%	34%	22%	4.9%	17%	27%	
30-44	32%	26%	39%	32%	5.7%	27%	38%	
45-64	30%	25%	34%	31%	4.5%	27%	36%	
65+	25%	20%	31%	28%	6.0%	22%	34%	
< High School	NSR			25%	8.7%	16%	33%	
High School	31%	26%	36%	32%	4.5%	28%	36%	
Some College	25%	19%	32%	35%	5.0%	30%	40%	
College Graduate	23%	18%	28%	19%	4.6%	15%	24%	
<\$25,000	32%	26%	39%	33%	5.3%	28%	38%	
\$25,000 to \$49,999	27%	22%	33%	25%	5.2%	20%	30%	
\$50,000+	29%	24%	34%	30%	4.6%	25%	34%	
White, non-Hispanic	27%	24%	31%	28%	2.7%	25%	31%	
Black, non-Hispanic	NA			38%	13.7%	24%	51%	
Hispanic	NA			35%	17.3%	17%	52%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q9.1-9.5: Percent of adults who eat fruits and vegetables 5+ times per day

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	21%	18%	24%	10%	1.7%	8%	11%	***
Male	15%	11%	20%	8%	2.2%	6%	10%	***
Female	25%	22%	29%	11%	2.6%	9%	14%	***
18-29	25%	17%	37%	10%	3.6%	6%	13%	***
30-44	16%	12%	22%	9%	3.4%	5%	12%	***
45-64	19%	15%	23%	10%	2.9%	7%	13%	***
65+	25%	20%	30%	10%	4.1%	5%	14%	***
< High School	NSR			6%	4.9%	1%	11%	
High School	17%	13%	21%	10%	2.8%	7%	12%	***
Some College	20%	15%	27%	9%	3.0%	6%	12%	***
College Graduate	29%	23%	35%	10%	3.6%	7%	14%	***
<\$25,000	20%	15%	27%	7%	2.9%	4%	10%	***
\$25,000 to \$49,999	18%	13%	23%	8%	3.3%	4%	11%	***
\$50,000+	24%	19%	29%	13%	3.3%	9%	16%	***
White, non-Hispanic	20%	17%	23%	10%	1.8%	8%	11%	***
Black, non-Hispanic	NA			10%	8.6%	2%	19%	
Hispanic	NA			14%	13.0%	1%	27%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q10.1: Percent of adults with no leisure time physical activity in past month

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	24%	21%	27%	28%	2.6%	25%	31%	
Male	19%	15%	24%	25%	3.5%	21%	28%	
Female	29%	25%	33%	31%	3.7%	28%	35%	
18-29	18%	11%	27%	24%	5.0%	19%	29%	
30-44	22%	17%	28%	28%	5.4%	23%	33%	
45-64	24%	20%	28%	28%	4.3%	24%	32%	
65+	35%	29%	41%	33%	6.2%	27%	39%	
< High School	40%	28%	53%	51%	10.6%	40%	61%	
High School	28%	24%	33%	36%	4.5%	31%	40%	
Some College	22%	17%	29%	30%	4.8%	25%	35%	
College Graduate	15%	11%	21%	13%	4.0%	9%	17%	
<\$25,000	39%	32%	47%	39%	5.4%	34%	44%	
\$25,000 to \$49,999	22%	17%	27%	24%	5.1%	19%	29%	
\$50,000+	14%	10%	19%	21%	4.0%	17%	25%	
White, non-Hispanic	24%	21%	27%	27%	2.6%	25%	30%	
Black, non-Hispanic	NA			29%	12.7%	16%	41%	
Hispanic	NA			48%	18.2%	30%	66%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q10.2: Percent of adults that participated in each activity* (Out of adults that participated in leisure time activities in past month)

Activity	%
Walking	53%
Running	10%
Gardening (spading, weeding, digging, filling)	5%
Weight lifting	4%
Other	4%
Aerobics video or class	3%
Bicycling	3%
Elliptical/EFX machine exercise	3%
Bicycling machine exercise	2%
Basketball	1%
Bowling	1%
Calisthenics	1%
Dancing-ballet, ballroom, Latin, hip hop, etc.	1%
Golf (with motorized cart)	1%
Golf (without motorized cart)	1%
Hunting large game-deer, elk	1%
Jogging	1%
Pilates	1%
Swimming	1%
Yoga	1%
Boxing	<1%
Fishing from river or boat	<1%
Hiking- cross country	<1%
Horseback riding	<1%
Hunting small game- quail	<1%
Mowing lawn	<1%
Raking lawn	<1%
Rowing machine exercise	<1%
Skateboarding	<1%
Snow shoveling by hand	<1%
Soccer	<1%
Stair climbing/Stair master	<1%
Swimming in laps	<1%
Tai Chi	<1%
Tennis	<1%
Touch football	<1%
Volleyball	<1%

*Respondents were asked ""What type of physical activity or exercise did you spend *the most* time doing during the past month?"

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q10.4: Average number of times spent exercising per week
(of adults who exercised in past month)

	2007			2011				Sig
	Avg #	Lower	Upper	%	Avg #	Lower	Upper	
Total	NA			NA	3.8	2.5	5.1	
Male	NA			NA	3.7	3.5	4.0	
Female	NA			NA	3.9	3.6	4.1	
18-29	NA			NA	3.7	3.3	4.0	
30-44	NA			NA	3.7	3.3	4.0	
45-64	NA			NA	3.9	3.7	4.2	
65+	NA			NA	3.7	3.3	4.2	
< High School	NA			NA	4.3	3.5	5.1	
High School	NA			NA	3.9	3.6	4.2	
Some College	NA			NA	3.8	3.4	4.1	
College Graduate	NA			NA	3.6	3.4	3.9	
<\$25,000	NA			NA	4.2	3.8	4.6	
\$25,000 to \$49,999	NA			NA	3.4	3.1	3.7	
\$50,000+	NA			NA	3.8	3.5	4.1	
White, non-Hispanic	NA			NA	3.8	3.6	3.9	
Black, non-Hispanic	NA			NA	3.9	2.8	4.9	
Hispanic	NA			NA	4.9	3.8	6.0	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q10.4: Average minutes per week spent exercising (of adults that exercised in past month)

	2007			2011				Sig
	%	Lower	Upper	%	Avg Min.	Lower	Upper	
Total	NA			NA	58.6	55.3	62.0	
Male	NA			NA	67.4	62.0	72.8	
Female	NA			NA	49.9	46.0	53.7	
18-29	NA			NA	59.7	54.2	65.1	
30-44	NA			NA	56.4	49.5	63.4	
45-64	NA			NA	61.7	55.3	68.0	
65+	NA			NA	53.7	45.4	62.1	
< High School	NA			NA	62.1	49.1	75.1	
High School	NA			NA	60.3	54.3	66.3	
Some College	NA			NA	58.7	51.6	65.7	
College Graduate	NA			NA	56.6	51.3	61.9	
<\$25,000	NA			NA	55.4	48.2	62.7	
\$25,000 to \$49,999	NA			NA	57.2	50.8	63.5	
\$50,000+	NA			NA	57.7	52.1	63.4	
White, non-Hispanic	NA			NA	59.1	55.6	62.7	
Black, non-Hispanic	NA			NA	42.2	32.8	51.7	
Hispanic	NA			NA	60.1	39.6	80.6	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q10.5: Average number of times spent strengthening muscles per week
(of adults that did muscle strength activities in past month)

	2007			2011				Sig
	%	Lower	Upper	%	Avg #.	Lower	Upper	
Total	NA			NA	1.0	0.5	1.6	
Male	NA			NA	1.2	1.1	1.4	
Female	NA			NA	0.9	0.7	1.0	
18-29	NA			NA	1.5	1.2	1.7	
30-44	NA			NA	1.1	0.9	1.4	
45-64	NA			NA	0.8	0.7	1.0	
65+	NA			NA	0.8	0.6	1.0	
< High School	NA			NA	0.7	0.3	1.0	
High School	NA			NA	0.8	0.7	1.0	
Some College	NA			NA	1.1	0.9	1.3	
College Graduate	NA			NA	1.3	1.1	1.5	
<\$25,000	NA			NA	0.9	0.6	1.1	
\$25,000 to \$49,999	NA			NA	0.9	0.7	1.1	
\$50,000+	NA			NA	1.4	1.2	1.6	
White, non-Hispanic	NA			NA	1.1	1.0	1.2	
Black, non-Hispanic	NA			NA	0.7	0.4	1.1	
Hispanic	NA			NA	0.9	0.3	1.4	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q11.1: Percent of adults with health problems that require use of special equipment

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	6%	4%	7%	8%	1.6%	7%	10%	***
Male	4%	3%	7%	8%	2.2%	6%	10%	
Female	7%	5%	9%	8%	2.2%	6%	10%	
18-29	1%	0%	8%	2%	1.7%	0%	4%	
30-44	2%	1%	6%	6%	2.9%	3%	9%	
45-64	6%	5%	9%	9%	2.7%	6%	12%	
65+	13%	9%	18%	17%	4.9%	12%	22%	
< High School	7%	3%	15%	22%	8.3%	13%	30%	
High School	6%	4%	8%	10%	2.8%	7%	13%	
Some College	6%	4%	10%	7%	2.7%	5%	10%	
College Graduate	5%	3%	8%	5%	2.5%	2%	7%	
<\$25,000	11%	7%	15%	18%	4.3%	14%	23%	
\$25,000 to \$49,999	4%	2%	6%	5%	2.6%	2%	8%	
\$50,000+	3%	1%	5%	4%	1.9%	2%	6%	
White, non-Hispanic	5%	4%	7%	8%	1.6%	6%	10%	
Black, non-Hispanic	NA			2%	4.1%	0%	6%	
Hispanic	NA			17%	13.7%	3%	31%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q12.1: Percent of adults who have arthritis or joint pain that limits activity

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			43%	5.1%	38%	48%	
Male	NA			49%	8.1%	41%	57%	
Female	NA			39%	6.6%	32%	45%	
18-29	NA			NSR				
30-44	NA			58%	15.3%	42%	73%	
45-64	NA			43%	7.4%	36%	50%	
65+	NA			37%	8.2%	29%	45%	
< High School	NA			58%	17.1%	41%	75%	
High School	NA			49%	7.9%	41%	56%	
Some College	NA			43%	9.5%	34%	53%	
College Graduate	NA			30%	11.6%	18%	41%	
<\$25,000	NA			49%	8.9%	40%	58%	
\$25,000 to \$49,999	NA			36%	10.8%	26%	47%	
\$50,000+	NA			38%	10.6%	28%	49%	
White, non-Hispanic	NA			42%	5.3%	37%	48%	
Black, non-Hispanic	NA			40%	30.4%	10%	70%	
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q13.1: Percent of adults who always wear a seat belt when they drive or ride in a car

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	74%	70%	77%	83%	2.2%	80%	85%	***
Male	67%	62%	73%	78%	3.4%	75%	82%	***
Female	80%	75%	83%	87%	2.7%	84%	89%	***
18-29	64%	52%	74%	77%	5.0%	72%	82%	
30-44	73%	66%	79%	79%	4.9%	74%	84%	
45-64	77%	72%	81%	85%	3.4%	82%	89%	***
65+	79%	73%	84%	89%	4.1%	85%	93%	***
< High School	NSR			93%	8.9%	84%	100%	
High School	73%	67%	78%	88%	3.9%	84%	92%	***
Some College	76%	69%	82%	79%	4.2%	75%	83%	
College Graduate	81%	75%	86%	72%	3.2%	69%	75%	***
<\$25,000	68%	60%	75%	77%	4.7%	73%	82%	
\$25,000 to \$49,999	77%	70%	82%	78%	4.0%	74%	82%	
\$50,000+	77%	72%	82%	92%	3.7%	89%	96%	***
White, non-Hispanic	74%	71%	77%	83%	2.2%	81%	85%	***
Black, non-Hispanic	NA			65%	13.3%	52%	79%	
Hispanic	NA			79%	14.7%	65%	94%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q14.1: Percent of adults with seasonal flu shot or flu vaccine that was sprayed up nose in past year, all ages

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			45%	2.8%	42%	48%	
Male	NA			45%	4.1%	41%	49%	
Female	NA			45%	3.9%	41%	49%	
18-29	NA			24%	5.1%	19%	29%	
30-44	NA			35%	5.7%	29%	40%	
45-64	NA			52%	4.8%	47%	56%	
65+	NA			70%	6.1%	64%	76%	
< High School	NA			39%	10.2%	29%	50%	
High School	NA			54%	4.8%	49%	58%	
Some College	NA			41%	5.2%	36%	46%	
College Graduate	NA			37%	5.3%	32%	42%	
<\$25,000	NA			40%	5.5%	35%	46%	
\$25,000 to \$49,999	NA			42%	5.7%	36%	47%	
\$50,000+	NA			52%	5.1%	47%	57%	
White, non-Hispanic	NA			45%	2.9%	42%	48%	
Black, non-Hispanic	NA			35%	13.3%	21%	48%	
Hispanic	NA			41%	17.9%	23%	59%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q14.1: Percent of adults with seasonal flu shot or flu vaccine that was sprayed up nose in past year, age 50+

	2007*			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	58%	54%	63%	60%	4.0%	55%	64%	
Male	55%	48%	61%	60%	5.8%	55%	66%	
Female	62%	56%	67%	59%	5.5%	53%	64%	
18-29	NA			NA				
30-49	NA			NA				
50-64	44%	39%	50%	53%	5.2%	48%	58%	
65+	76%	70%	82%	70%	6.1%	64%	76%	
< High School	NSR			60%	15.1%	45%	75%	
High School	61%	55%	67%	66%	6.2%	60%	72%	
Some College	58%	49%	67%	57%	8.1%	49%	65%	
College Graduate	55%	47%	63%	51%	7.9%	43%	58%	
<\$25,000	64%	56%	72%	51%	7.9%	43%	59%	
\$25,000 to \$49,999	58%	51%	66%	60%	8.0%	52%	68%	
\$50,000+	54%	46%	61%	62%	7.5%	55%	70%	
White, non-Hispanic	59%	54%	63%	60%	4.1%	56%	64%	
Black, non-Hispanic	NA			53%	25.2%	28%	79%	
Hispanic	NA			NSR				

*2007 results do not include adults receiving the *nasal spray* vaccine. Because only 1% of adults received the vaccine in the form of a spray in 2007, the data is still comparable.

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q14.1: Percent of adults with seasonal flu shot or flu vaccine that was sprayed up nose in past year, age 65+

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	76%	70%	82%	70%	6.1%	64%	76%	
Male	75%	64%	83%	73%	9.0%	64%	82%	
Female	78%	69%	84%	68%	8.2%	59%	76%	
18-29	NA			NA				
30-44	NA			NA				
45-64	NA			NA				
65+	76%	70%	82%	70%	6.1%	64%	76%	
< High School	NSR			70%	19.7%	50%	90%	
High School	80%	72%	87%	80%	8.7%	71%	88%	
Some College	NSR			60%	13.3%	46%	73%	
College Graduate	76%	64%	85%	66%	12.0%	54%	78%	
<\$25,000	75%	64%	84%	64%	10.9%	53%	75%	
\$25,000 to \$49,999	77%	66%	85%	70%	11.5%	59%	82%	
\$50,000+	NSR			73%	16.1%	57%	89%	
White, non-Hispanic	77%	71%	82%	70%	6.1%	64%	76%	
Black, non-Hispanic	NA			NSR				
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q14.3: Place where received last flu shot/nasal vaccine spray

Location of Last Vaccination	%
A doctors office or health maintenance organization (HMO)	40%
Workplace	24%
A store (EX: supermarket, drug store)	16%
A hospital	8%
A senior, recreation or community center	4%
Another type of clinic or health center	3%
Some other kind of place	3%
A health department	1%
A school	1%
An emergency room	<1%

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q14.4: Percent of adults who ever had a pneumonia vaccination, age 65+

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	71%	65%	77%	73%	6.0%	67%	79%	
Male	61%	50%	70%	74%	9.3%	64%	83%	
Female	78%	71%	84%	72%	7.9%	64%	80%	
18-29	NA			NA				
30-44	NA			NA				
45-64	NA			NA				
65+	71%	65%	77%	73%	6.0%	67%	79%	
< High School	NSR			96%	17.5%	78%	100%	
High School	73%	64%	81%	82%	8.6%	74%	91%	
Some College	NSR			67%	12.8%	55%	80%	
College Graduate	76%	64%	85%	59%	14.5%	45%	74%	
<\$25,000	77%	66%	85%	78%	9.7%	69%	88%	
\$25,000 to \$49,999	65%	54%	75%	66%	12.4%	54%	79%	
\$50,000+	NSR			65%	17.0%	48%	82%	
White, non-Hispanic	71%	65%	77%	73%	6.0%	67%	79%	
Black, non-Hispanic	NA			NSR				
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q15.1: Alcohol Consumption: Percent of adults who participated in Chronic Drinking in the past month

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	6%	4%	9%	6%	1.3%	4%	7%	
Male	10%	7%	15%	11%	2.6%	8%	13%	
Female	2%	1%	4%	1%	0.6%	0%	1%	***
18-29	12%	5%	24%	5%	2.5%	2%	7%	
30-44	6%	3%	11%	6%	2.9%	3%	9%	
45-64	5%	3%	7%	7%	2.4%	4%	9%	
65+	4%	2%	7%	4%	2.6%	1%	7%	
< High School	4%	1%	11%	3%	3.3%	0%	6%	
High School	7%	4%	11%	6%	2.2%	4%	8%	
Some College	6%	2%	15%	6%	2.5%	4%	9%	
College Graduate	6%	3%	11%	5%	2.6%	2%	8%	
<\$25,000	5%	2%	9%	6%	2.7%	3%	9%	
\$25,000 to \$49,999	6%	3%	13%	3%	2.2%	1%	6%	
\$50,000+	8%	5%	13%	9%	2.8%	6%	12%	
White, non-Hispanic	6%	4%	9%	6%	1.4%	4%	7%	
Black, non-Hispanic	NA			4%	5.7%	0%	10%	
Hispanic	NA			10%	11.1%	0%	21%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q15.2: Alcohol Consumption: Percent of adults who are heavy drinkers

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	6%	4%	8%	6%	1.3%	4%	7%	
Male	7%	4%	12%	9%	2.3%	6%	11%	
Female	4%	3%	6%	3%	1.4%	2%	4%	
18-29	13%	7%	25%	8%	3.2%	5%	11%	
30-44	5%	3%	8%	6%	2.9%	3%	9%	
45-64	4%	2%	6%	6%	2.2%	3%	8%	
65+	4%	2%	7%	3%	2.1%	1%	5%	
< High School	2%	1%	10%	3%	3.3%	0%	6%	
High School	6%	4%	10%	6%	2.3%	4%	9%	
Some College	6%	2%	15%	6%	2.4%	3%	8%	
College Graduate	6%	3%	11%	5%	2.7%	3%	8%	
<\$25,000	3%	2%	6%	6%	2.7%	4%	9%	
\$25,000 to \$49,999	6%	3%	13%	5%	2.6%	2%	7%	
\$50,000+	7%	4%	12%	7%	2.6%	5%	10%	
White, non-Hispanic	6%	4%	8%	6%	1.4%	4%	7%	
Black, non-Hispanic	NA			6%	6.9%	0%	13%	
Hispanic	NA			10%	11.1%	0%	21%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q15.3: Percent of adults who are binge drinkers

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	21%	18%	25%	19%	2.2%	17%	21%	
Male	29%	24%	35%	25%	3.6%	22%	29%	
Female	14%	12%	18%	13%	2.7%	10%	16%	
18-29	43%	32%	55%	31%	5.5%	25%	36%	
30-44	29%	23%	35%	24%	5.2%	19%	29%	
45-64	13%	10%	17%	15%	3.4%	11%	18%	
65+	6%	3%	10%	6%	3.0%	2%	9%	
< High School	NSR			10%	6.1%	4%	16%	
High School	17%	13%	22%	17%	3.5%	13%	20%	
Some College	28%	21%	36%	19%	4.1%	15%	23%	
College Graduate	21%	16%	27%	20%	4.7%	16%	25%	
<\$25,000	22%	15%	29%	15%	4.1%	11%	19%	
\$25,000 to \$49,999	23%	17%	30%	16%	4.4%	11%	20%	
\$50,000+	24%	19%	30%	31%	4.6%	27%	36%	
White, non-Hispanic	22%	19%	25%	19%	2.3%	16%	21%	
Black, non-Hispanic	NA			28%	12.8%	15%	40%	
Hispanic	NA			19%	14.6%	4%	33%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q16.1: Percent of adults ever tested for HIV, age 18-64

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	29%	26%	33%	40%	3.1%	36%	43%	***
Male	25%	20%	31%	36%	4.3%	32%	40%	***
Female	34%	29%	39%	43%	4.5%	39%	48%	***
18-29	27%	19%	37%	41%	5.9%	35%	47%	
30-44	47%	40%	54%	52%	6.0%	46%	58%	
45-64	17%	14%	21%	31%	4.5%	26%	35%	***
65+	NA			NA				
< High School	NSR			61%	13.0%	48%	74%	
High School	25%	20%	32%	40%	5.4%	34%	45%	***
Some College	35%	27%	43%	44%	5.7%	38%	50%	
College Graduate	28%	22%	35%	29%	5.6%	24%	35%	
<\$25,000	34%	26%	44%	52%	6.5%	45%	59%	***
\$25,000 to \$49,999	30%	23%	37%	38%	6.3%	32%	45%	
\$50,000+	28%	22%	34%	37%	5.1%	32%	42%	
White, non-Hispanic	29%	25%	33%	37%	3.2%	34%	40%	***
Black, non-Hispanic	NA			69%	13.1%	56%	82%	
Hispanic	NA			61%	18.1%	43%	79%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Q16.2: Percent of adults who practiced risky behavior within the past year that could result in HIV infection

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			2%	0.8%	1%	3%	
Male	NA			2%	1.2%	1%	3%	
Female	NA			2%	1.1%	1%	3%	
18-29	NA			2%	1.6%	0%	3%	
30-44	NA			4%	2.4%	2%	7%	
45-64	NA			2%	1.2%	0%	3%	
65+	NA			1%	1.0%	0%	1%	
< High School	NA			0%	NCI			
High School	NA			3%	1.5%	1%	4%	
Some College	NA			2%	1.4%	0%	3%	
College Graduate	NA			2%	1.5%	0%	3%	
<\$25,000	NA			3%	1.8%	1%	4%	
\$25,000 to \$49,999	NA			1%	1.0%	0%	2%	
\$50,000+	NA			2%	1.4%	1%	3%	
White, non-Hispanic	NA			2%	0.9%	1%	3%	
Black, non-Hispanic	NA			2%	3.9%	0%	6%	
Hispanic	NA			0%	NCI			

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 1, 1: Percent of adults who have ever been told they have pre-diabetes

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	NA			6%	1.5%	5%	8%	
Male	NA			8%	2.3%	5%	10%	
Female	NA			5%	1.9%	3%	7%	
18-29	NA			3%	2.1%	1%	5%	
30-44	NA			3%	2.0%	1%	5%	
45-64	NA			9%	2.9%	6%	12%	
65+	NA			11%	4.7%	7%	16%	
< High School	NA			6%	5.1%	1%	11%	
High School	NA			8%	2.7%	6%	11%	
Some College	NA			6%	2.7%	4%	9%	
College Graduate	NA			4%	2.5%	2%	7%	
<\$25,000	NA			8%	3.3%	5%	11%	
\$25,000 to \$49,999	NA			6%	3.0%	3%	9%	
\$50,000+	NA			5%	2.3%	3%	8%	
White, non-Hispanic	NA			7%	1.5%	5%	8%	
Black, non-Hispanic	NA			2%	4.3%	0%	6%	
Hispanic	NA			11%	11.8%	0%	23%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 7, 1: Percent of adults who sleep less than 7 hours in a 24 hour period

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			37%	2.8%	34%	39%	
Male	NA			37%	3.9%	33%	41%	
Female	NA			37%	3.8%	33%	41%	
18-29	NA			38%	5.8%	32%	44%	
30-44	NA			44%	5.9%	38%	50%	
45-64	NA			34%	4.5%	30%	39%	
65+	NA			30%	6.2%	24%	37%	
< High School	NA			41%	10.5%	31%	52%	
High School	NA			41%	4.7%	36%	46%	
Some College	NA			39%	5.1%	34%	44%	
College Graduate	NA			28%	5.1%	23%	33%	
<\$25,000	NA			40%	5.5%	34%	45%	
\$25,000 to \$49,999	NA			32%	5.5%	26%	37%	
\$50,000+	NA			42%	4.9%	37%	47%	
White, non-Hispanic	NA			37%	2.9%	34%	40%	
Black, non-Hispanic	NA			27%	12.4%	14%	39%	
Hispanic	NA			32%	17.3%	15%	49%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module Erie HC, 1: Place where usually go when sick

Location	%
Doctors office	71%
No usual place	15%
A hospital emergency room	5%
Urgent care center	4%
A hospital outpatient department	3%
A public health clinic or community health center	2%
Some other kind of place	1%

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module Erie HC, 2: Where went for care when sick in past 12 months

Location	%
Doctors office	64%
No usual place	23%
A hospital emergency room	9%
Urgent care center	5%
A hospital outpatient department	4%
A public health clinic or community health center	2%
Some other kind of place	2%

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 12, 1: Percent of women who had a mammogram in the past year, women age 40+

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	65%	61%	70%	67%	4.8%	62%	72%	
Male	NA			NA				
Female	65%	61%	70%	67%	4.8%	62%	72%	
18-39	NA			NA				
40-44	NA			50%	15.9%	34%	66%	
45-54	NA			65%	6.4%	58%	71%	
65+	NA			77%	7.4%	69%	84%	
< High School	NSR			94%	15.2%	79%	100%	
High School	66%	60%	72%	72%	7.4%	65%	80%	
Some College	65%	55%	74%	63%	9.5%	53%	72%	
College Graduate	71%	62%	79%	58%	9.3%	48%	67%	
<\$25,000	57%	48%	66%	54%	9.2%	45%	63%	
\$25,000 to \$49,999	71%	62%	78%	66%	9.6%	57%	76%	
\$50,000+	72%	64%	79%	80%	8.2%	72%	88%	
White, non-Hispanic	65%	60%	70%	67%	4.9%	62%	72%	
Black, non-Hispanic	NA			69%	25.1%	44%	94%	
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 12, 1: Percent of women who never had a mammogram, women age 40+

	2007		2011				Sig
	%	Lower Upper	%	CI	Lower	Upper	
Total	5%	NA	6%	2.4%	4%	8%	
Male	NA	NA	NA				
Female	5%	NA	6%	2.4%	4%	8%	
18-39	NA	NA	NA				
40-44	NA	NA	24%	13.5%	10%	37%	
45-54	NA	NA	4%	2.7%	2%	7%	
65+	NA	NA	3%	3.1%	0%	6%	
< High School	6%	NA	0%	NCI			
High School	5%	NA	8%	4.0%	4%	12%	
Some College	3%	NA	7%	5.1%	2%	12%	
College Graduate	7%	NA	4%	4.3%	0%	8%	
<\$25,000	6%	NA	4%	3.8%	1%	8%	
\$25,000 to \$49,999	5%	NA	2%	3.4%	0%	6%	
\$50,000+	3%	NA	9%	5.0%	4%	14%	
White, non-Hispanic	5%	NA	6%	2.5%	4%	9%	
Black, non-Hispanic	NA	NA	0%	NCI			
Hispanic	NA	NA	NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 12, 2: Percent of women who ever had a clinical breast exam, women age 40+

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	93%	90%	95%	97%	1.9%	95%	98%	***
Male	NA			NA%				
Female	93%	90%	95%	97%	1.9%	95%	98%	***
18-39	NA			NA				
40-44	NA			95%	7.1%	88%	100%	
45-54	NA			97%	2.4%	94%	99%	
65+	NA			97%	3.2%	94%	100%	
< High School	NSR			100%	NCI			
High School	92%	88%	95%	100%	3.7%	96%	100%	
Some College	96%	88%	99%	99%	2.0%	97%	100%	
College Graduate	95%	88%	98%	78%	3.1%	75%	81%	***
<\$25,000	89%	82%	94%	95%	4.2%	90%	99%	
\$25,000 to \$49,999	94%	88%	97%	89%	2.4%	87%	91%	
\$50,000+	98%	93%	99%	100%	3.5%	97%	100%	
White, non-Hispanic	93%	90%	95%	97%	1.9%	95%	99%	***
Black, non-Hispanic	NA			100%	NCI			
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 12, 2: Percent of women who had a clinical breast exam in the past year, women age 40+

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	64%	NA		67%	4.8%	63%	72%	
Male	NA			NA				
Female	64%	NA		67%	4.8%	63%	72%	
18-39	NA			NA				
40-44	NA			66%	15.1%	51%	81%	
45-64	NA			68%	6.3%	62%	74%	
65+	NA			66%	8.4%	58%	75%	
< High School	36%	NA		84%	17.2%	67%	100%	
High School	65%	NA		72%	7.5%	65%	80%	
Some College	63%	NA		64%	9.5%	55%	74%	
College Graduate	71%	NA		58%	9.2%	49%	68%	
<\$25,000	55%	NA		55%	9.3%	46%	64%	
\$25,000 to \$49,999	70%	NA		65%	9.9%	55%	75%	
\$50,000+	72%	NA		83%	8.0%	75%	91%	
White, non-Hispanic	64%	NA		67%	5.0%	62%	72%	
Black, non-Hispanic	NA			77%	22.9%	54%	100%	
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 12, 3: Percent of women who had a pap test in the past year, women age 18+

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	59%	54%	63%	60%	3.9%	56%	64%	
Male	NA			NA				
Female	59%	54%	63%	60%	3.9%	56%	64%	
18-29	NSR			66%	8.0%	58%	74%	
30-44	64%	55%	72%	74%	7.4%	67%	81%	
45-64	62%	56%	68%	61%	6.6%	55%	68%	
65+	31%	24%	39%	33%	8.4%	24%	41%	
< High School	NSR			47%	14.6%	33%	62%	
High School	54%	48%	61%	58%	6.6%	51%	65%	
Some College	60%	51%	68%	60%	7.3%	53%	67%	
College Graduate	69%	61%	76%	60%	6.7%	53%	67%	
<\$25,000	45%	36%	54%	48%	7.4%	41%	55%	
\$25,000 to \$49,999	65%	57%	71%	55%	7.8%	48%	63%	
\$50,000+	70%	62%	76%	81%	6.5%	75%	88%	
White, non-Hispanic	59%	54%	63%	58%	4.1%	54%	62%	
Black, non-Hispanic	NA			93%	9.5%	83%	100%	
Hispanic	NA			54%	27.1%	27%	81%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 13, 1: Percent of men who had a prostate-specific antigen test in the past year, men age 40+

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	44%	38%	50%	52%	5.5%	46%	57%	
Male	44%	38%	50%	52%	5.5%	46%	57%	
Female	NA			NA				
18-39	NA			NA				
40-44	NA			15%	11.9%	3%	27%	
45-64	NA			49%	7.0%	42%	56%	
65+	NA			72%	9.3%	63%	81%	
< High School	NSR			57%	23.7%	33%	80%	
High School	41%	32%	51%	55%	8.9%	46%	64%	
Some College	50%	37%	63%	46%	10.4%	35%	56%	
College Graduate	44%	34%	54%	48%	9.8%	39%	58%	
<\$25,000	NSR			47%	11.9%	35%	59%	
\$25,000 to \$49,999	52%	42%	63%	44%	11.0%	33%	55%	
\$50,000+	41%	33%	51%	60%	9.1%	51%	69%	***
White, non-Hispanic	44%	38%	50%	51%	5.6%	46%	57%	
Black, non-Hispanic	NA			NSR				
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 13, 2: Percent of men who had a digital rectal exam in the past year, men age 40+

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	46%	40%	52%	42%	5.3%	37%	47%	
Male	46%	40%	52%	42%	5.3%	37%	47%	
Female	NA			NA				
18-39	NA			NA				
40-44	NA			12%	10.8%	1%	23%	
45-64	NA			42%	6.7%	35%	48%	
65+	NA			55%	10.3%	45%	65%	
< High School	NSR			23%	19.1%	3%	42%	
High School	44%	34%	53%	44%	8.7%	35%	52%	
Some College	51%	38%	64%	40%	10.0%	30%	50%	
College Graduate	46%	36%	56%	40%	9.9%	30%	50%	
<\$25,000	NSR			39%	11.8%	27%	50%	
\$25,000 to \$49,999	45%	35%	56%	29%	10.1%	18%	39%	
\$50,000+	47%	38%	56%	51%	9.1%	42%	60%	
White, non-Hispanic	46%	40%	52%	43%	10.8%	32%	53%	
Black, non-Hispanic	NA			NSR				
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 13, 3: Percent of men who have ever been told they had prostate cancer, men age 40+

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	5%	3%	8%	6%	2.5%	4%	9%	
Male	5%	3%	8%	6%	2.5%	4%	9%	
Female	NA			NA				
18-39	NA			NA				
40-44	NA			0%	NCI			
45-64	NA			3%	2.4%	1%	6%	
65+	NA			15%	7.2%	8%	22%	
< High School	NSR			24%	20.2%	4%	44%	
High School	5%	2%	10%	7%	4.2%	3%	11%	
Some College	5%	2%	14%	3%	3.5%	0%	7%	
College Graduate	4%	2%	10%	6%	5.0%	1%	11%	
<\$25,000	2%	1%	9%	11%	7.4%	4%	19%	
\$25,000 to \$49,999	11%	6%	18%	4%	4.6%	0%	9%	
\$50,000+	3%	1%	8%	6%	4.2%	2%	11%	
White, non-Hispanic	5%	3%	8%	6%	2.5%	3%	8%	
Black, non-Hispanic	NA			NSR				
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 14, 1: Percent of adults who had a blood stool test in the past 2 years, age 50+

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	25%	22%	29%	18%	3.2%	15%	21%	***
Male	30%	24%	36%	19%	4.8%	15%	24%	***
Female	22%	18%	27%	16%	4.2%	12%	20%	
18-29	NA			NA				
30-49	NA			NA				
50-64	NA			14%	3.7%	11%	18%	
65+	NA			24%	5.7%	18%	29%	
< High School	NSR			35%	14.4%	20%	49%	
High School	25%	20%	32%	17%	4.6%	13%	22%	
Some College	23%	16%	32%	15%	5.9%	9%	21%	
College Graduate	28%	22%	36%	16%	6.6%	9%	23%	
<\$25,000	26%	19%	35%	20%	6.4%	13%	26%	
\$25,000 to \$49,999	31%	24%	38%	16%	6.5%	9%	22%	***
\$50,000+	24%	18%	31%	21%	6.0%	15%	27%	
White, non-Hispanic	24%	21%	28%	18%	3.2%	14%	21%	***
Black, non-Hispanic	NA			20%	20.2%	0%	40%	
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 14, 2: Percent of adults who had a sigmoidoscopy or colonoscopy in the past 5 years, age 50+

	2007			2011				Sig
	%	Lower	Upper	%	CI	Lower	Upper	
Total	53%	49%	58%	62%	4.0%	58%	66%	***
Male	56%	49%	63%	66%	5.7%	60%	71%	
Female	51%	45%	56%	60%	5.6%	54%	65%	
18-29	NA			NA				
30-49	NA			NA				
50-64	NA			60%	5.1%	55%	65%	
65+	NA			67%	6.3%	60%	73%	
< High School	NSR			67%	15.2%	52%	83%	
High School	49%	43%	55%	67%	6.2%	61%	74%	***
Some College	53%	43%	62%	60%	8.0%	52%	68%	
College Graduate	65%	57%	72%	55%	7.6%	47%	63%	
<\$25,000	48%	40%	57%	54%	7.9%	46%	62%	
\$25,000 to \$49,999	54%	46%	61%	55%	8.3%	47%	64%	
\$50,000+	56%	49%	64%	74%	7.2%	67%	82%	***
White, non-Hispanic	53%	49%	58%	62%	4.1%	58%	66%	***
Black, non-Hispanic	NA			64%	25.1%	39%	89%	
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 23, 1: Percent of adults who feel household is well prepared to handle a large scale disaster or emergency

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	19%	16%	22%	28%	2.6%	25%	30%	***
Male	25%	20%	31%	34%	3.9%	30%	38%	
Female	14%	11%	17%	22%	3.3%	18%	25%	***
18-29	24%	14%	36%	24%	5.1%	19%	29%	
30-44	13%	9%	18%	24%	5.1%	19%	29%	***
45-64	18%	14%	22%	29%	4.3%	24%	33%	***
65+	26%	21%	33%	36%	6.5%	29%	42%	
< High School	NSR			32%	9.9%	22%	41%	
High School	22%	17%	28%	31%	4.4%	27%	35%	
Some College	21%	14%	29%	28%	4.7%	23%	33%	
College Graduate	12%	9%	17%	22%	4.8%	17%	27%	***
<\$25,000	23%	16%	31%	27%	5.0%	22%	32%	
\$25,000 to \$49,999	17%	12%	23%	22%	5.0%	17%	27%	
\$50,000+	15%	11%	19%	31%	4.6%	27%	36%	***
White, non-Hispanic	19%	16%	22%	29%	2.7%	26%	31%	***
Black, non-Hispanic	NA			18%	10.8%	8%	29%	
Hispanic	NA			11%	11.8%	0%	23%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 28, 1: Percent of adults who were worried or stressed about having enough money

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			32%	2.8%	29%	35%	
Male	NA			26%	3.8%	23%	30%	
Female	NA			37%	3.9%	33%	41%	
18-29	NA			47%	6.7%	40%	54%	
30-44	NA			43%	6.0%	37%	49%	
45-64	NA			26%	4.2%	22%	30%	
65+	NA			15%	4.8%	10%	20%	
< High School	NA			46%	11.4%	35%	58%	
High School	NA			39%	4.8%	34%	44%	
Some College	NA			37%	5.2%	32%	42%	
College Graduate	NA			17%	4.5%	13%	22%	
<\$25,000	NA			53%	5.8%	47%	59%	
\$25,000 to \$49,999	NA			29%	5.6%	23%	34%	
\$50,000+	NA			20%	4.1%	16%	25%	
White, non-Hispanic	NA			31%	2.8%	28%	34%	
Black, non-Hispanic	NA			44%	14.5%	30%	59%	
Hispanic	NA			35%	18.3%	16%	53%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 28, 2: Percent of adults who needed prescribed medication in past 12 months but did not get it due to cost

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			12%	1.8%	10%	14%	
Male	NA			8%	2.2%	6%	10%	
Female	NA			16%	2.9%	13%	19%	
18-29	NA			16%	4.4%	12%	20%	
30-44	NA			15%	4.2%	11%	19%	
45-64	NA			11%	3.0%	8%	14%	
65+	NA			4%	2.6%	1%	7%	
< High School	NA			19%	7.8%	11%	27%	
High School	NA			15%	3.3%	11%	18%	
Some College	NA			13%	3.6%	10%	17%	
College Graduate	NA			6%	2.9%	4%	9%	
<\$25,000	NA			22%	4.7%	18%	27%	
\$25,000 to \$49,999	NA			11%	3.7%	7%	15%	
\$50,000+	NA			6%	2.3%	4%	8%	
White, non-Hispanic	NA			12%	1.9%	10%	13%	
Black, non-Hispanic	NA			14%	9.8%	4%	24%	
Hispanic	NA			14%	12.6%	1%	26%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module Erie Tobacco, 1: Percent of adult smokers who smoke 1+ packs of cigarettes (20 or more cigarettes) per day (out of everyday smokers)

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	48%	39%	57%	39%	6.7%	32%	46%	
Male	NSR			42%	9.9%	32%	52%	
Female	48%	38%	58%	37%	9.2%	28%	46%	
18-29	NSR			28%	12.4%	16%	40%	
30-44	43%	30%	57%	39%	12.6%	26%	51%	
45-64	46%	35%	57%	45%	11.2%	34%	56%	
65+	NSR			47%	22.5%	25%	70%	
< High School	NSR			49%	20.5%	29%	70%	
High School	49%	37%	61%	48%	10.2%	38%	59%	
Some College	NSR			33%	11.5%	21%	44%	
College Graduate	NSR			30%	19.4%	11%	49%	
<\$25,000	NSR			40%	10.3%	29%	50%	
\$25,000 to \$49,999	44%	32%	58%	37%	13.8%	23%	50%	
\$50,000+	NSR			52%	17.3%	34%	69%	
White, non-Hispanic	49%	40%	58%	40%	7.2%	33%	47%	
Black, non-Hispanic	NA			20%	24.8%	0%	45%	
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 16, 1: Percent of adults where smoking is never allowed inside the house

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	68%	64%	71%	43%	5.9%	37%	49%	***
Male	68%	62%	73%	46%	8.7%	37%	55%	***
Female	67%	63%	71%	39%	7.9%	31%	47%	***
18-29	65%	53%	75%	54%	11.4%	43%	65%	
30-44	67%	60%	74%	47%	11.2%	36%	59%	***
45-64	69%	64%	73%	31%	9.1%	21%	40%	***
65+	69%	62%	74%	39%	19.9%	19%	59%	***
< High School	NSR			41%	17.3%	24%	59%	
High School	61%	55%	66%	41%	8.9%	33%	50%	***
Some College	67%	59%	74%	44%	10.4%	33%	54%	***
College Graduate	82%	77%	86%	45%	14.6%	31%	60%	***
<\$25,000	60%	52%	67%	34%	8.8%	25%	43%	***
\$25,000 to \$49,999	60%	54%	67%	37%	11.6%	25%	48%	***
\$50,000+	81%	76%	85%	73%	12.9%	60%	86%	
White, non-Hispanic	68%	64%	71%	43%	6.3%	36%	49%	***
Black, non-Hispanic	NA			24%	20.2%	3%	44%	
Hispanic	NA			NSR				

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module 26, 1: Percent of adults who have ever been told have an anxiety disorder

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			17%	2.1%	15%	19%	
Male	NA			13%	2.8%	11%	16%	
Female	NA			21%	3.2%	17%	24%	
18-29	NA			25%	5.1%	20%	30%	
30-44	NA			17%	4.5%	12%	21%	
45-64	NA			16%	3.5%	12%	19%	
65+	NA			10%	3.9%	6%	13%	
< High School	NA			22%	8.5%	14%	31%	
High School	NA			20%	3.7%	16%	23%	
Some College	NA			18%	4.0%	14%	22%	
College Graduate	NA			12%	3.8%	8%	16%	
<\$25,000	NA			30%	5.1%	24%	35%	
\$25,000 to \$49,999	NA			15%	4.3%	11%	19%	
\$50,000+	NA			11%	3.1%	8%	14%	
White, non-Hispanic	NA			17%	2.2%	15%	20%	
Black, non-Hispanic	NA			10%	8.5%	2%	19%	
Hispanic	NA			17%	13.3%	3%	30%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module Erie Alcohol Consumption: Percent of adults who drove 1 or more times when they perhaps had too much to drink in the past 30 days

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	6%	4%	8%	4%	1.1%	3%	5%	
Male	10%	7%	15%	6%	2.0%	4%	8%	
Female	2%	1%	4%	2%	1.1%	1%	3%	
18-29	14%	7%	25%	7%	3.1%	4%	10%	
30-44	9%	5%	14%	8%	3.2%	5%	11%	
45-64	3%	1%	5%	2%	1.3%	1%	3%	
65+	0%	0%	3%	0%	0.8%	0%	1%	
< High School	NSR			3%	3.3%	0%	6%	
High School	4%	2%	8%	4%	1.9%	2%	6%	
Some College	8%	4%	14%	5%	2.3%	3%	7%	
College Graduate	8%	4%	13%	3%	2.1%	1%	5%	
<\$25,000	5%	2%	10%	5%	2.3%	2%	7%	
\$25,000 to \$49,999	5%	3%	9%	3%	2.0%	1%	5%	
\$50,000+	8%	5%	14%	6%	2.2%	3%	8%	
White, non-Hispanic	6%	4%	9%	4%	1.2%	3%	5%	
Black, non-Hispanic	NA			2%	3.9%	0%	6%	
Hispanic	NA			10%	11.1%	0%	21%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module Erie Nutri, 1: Percent of adults who have 2 or more servings of dairy daily

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	69%	65%	73%	39%	2.8%	36%	41%	***
Male	67%	60%	73%	34%	3.9%	30%	38%	***
Female	71%	66%	75%	43%	3.9%	39%	47%	***
18-29	77%	64%	86%	45%	5.9%	39%	51%	***
30-44	74%	66%	81%	42%	5.9%	36%	47%	***
45-64	62%	56%	67%	37%	4.6%	33%	42%	***
65+	67%	58%	74%	29%	6.1%	23%	35%	***
< High School	NSR			39%	10.1%	29%	49%	
High School	68%	61%	73%	38%	4.6%	33%	42%	***
Some College	70%	61%	78%	39%	5.1%	34%	44%	***
College Graduate	74%	68%	80%	36%	5.3%	31%	42%	***
<\$25,000	62%	52%	71%	36%	5.4%	31%	41%	***
\$25,000 to \$49,999	71%	64%	78%	36%	5.6%	30%	42%	***
\$50,000+	72%	66%	77%	47%	5.0%	42%	52%	***
White, non-Hispanic	69%	65%	73%	40%	2.9%	37%	43%	***
Black, non-Hispanic	NA			27%	12.4%	14%	39%	
Hispanic	NA			21%	14.7%	6%	35%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module Erie Nutri, 2: Percent of adults who have 1 or more servings of grains daily

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	82%	79%	85%	64%	2.8%	62%	67%	***
Male	81%	76%	86%	62%	4.0%	58%	66%	***
Female	83%	80%	86%	66%	3.8%	63%	70%	***
18-29	86%	75%	92%	65%	5.7%	59%	70%	***
30-44	87%	82%	91%	61%	5.8%	56%	67%	***
45-64	77%	73%	81%	63%	4.7%	58%	67%	***
65+	81%	76%	86%	71%	6.1%	64%	77%	
< High School	80%	69%	88%	64%	10.8%	54%	75%	
High School	82%	77%	85%	70%	4.7%	65%	75%	***
Some College	80%	73%	86%	64%	5.1%	59%	69%	***
College Graduate	85%	81%	89%	54%	5.0%	49%	59%	***
<\$25,000	82%	76%	87%	61%	5.5%	55%	66%	***
\$25,000 to \$49,999	78%	71%	83%	57%	5.6%	51%	63%	***
\$50,000+	86%	82%	89%	74%	4.8%	69%	78%	***
White, non-Hispanic	82%	79%	85%	65%	2.9%	62%	68%	***
Black, non-Hispanic	NA			61%	13.6%	48%	75%	
Hispanic	NA			40%	17.5%	22%	58%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module Erie Nutri, 3: Average number of sugar sweetened beverages per day

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			6%	1.4%	5%	8%	
Male	NA			7%	2.2%	5%	10%	
Female	NA			5%	1.8%	4%	7%	
18-29	NA			8%	3.2%	5%	11%	
30-44	NA			11%	3.8%	7%	15%	
45-64	NA			5%	2.0%	3%	7%	
65+	NA			1%	1.6%	0%	3%	
< High School	NA			6%	4.7%	1%	11%	
High School	NA			7%	2.3%	4%	9%	
Some College	NA			10%	3.1%	7%	13%	
College Graduate	NA			3%	2.0%	1%	5%	
<\$25,000	NA			8%	3.1%	5%	11%	
\$25,000 to \$49,999	NA			7%	3.2%	4%	10%	
\$50,000+	NA			5%	2.2%	3%	8%	
White, non-Hispanic	NA			6%	1.4%	4%	7%	
Black, non-Hispanic	NA			14%	9.8%	4%	24%	
Hispanic	NA			18%	14.2%	4%	32%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.

2011 Behavioral Risk Factor Surveillance Survey
Erie County, Pennsylvania

Module Erie Oral, 1: Percent of adults who visited a dentist within the past year

	2007			2011				Sig
	%	Lower	Upper	%	C I	Lower	Upper	
Total	NA			70%	2.6%	67%	72%	
Male	NA			66%	3.9%	63%	70%	
Female	NA			73%	3.5%	69%	76%	
18-29	NA			73%	5.3%	68%	78%	
30-44	NA			72%	5.4%	67%	77%	
45-64	NA			72%	4.3%	68%	77%	
65+	NA			59%	6.5%	52%	65%	
< High School	NA			61%	10.8%	50%	71%	
High School	NA			68%	4.7%	63%	72%	
Some College	NA			72%	4.7%	68%	77%	
College Graduate	NA			66%	4.1%	62%	70%	
<\$25,000	NA			53%	5.6%	47%	59%	
\$25,000 to \$49,999	NA			64%	5.2%	59%	69%	
\$50,000+	NA			92%	3.8%	88%	95%	
White, non-Hispanic	NA			70%	2.7%	67%	73%	
Black, non-Hispanic	NA			68%	12.9%	55%	81%	
Hispanic	NA			76%	15.6%	60%	91%	

NSR indicates the total response is less than 10 and the percentage is considered "not statistically reliable".

NCI indicates the percentage was 0.0% or 100.0% and no confidence interval was calculated.

NA indicates the necessary data is not available

Note: Significance tests compare 2007 data to 2011.